


**RESTRICTED USE PESTICIDE  
DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS**

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION  
AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.



**Adjourn™**  
INSECTICIDE

**ACTIVE INGREDIENT**

**BY WEIGHT**

Esfenvalerate	
(S)-cyano (3-phenoxyphenyl) methyl	
(S)-4-chloro-alpha-(1-methylethyl)benzeneacetate .....	8.4%
<b>INERT INGREDIENTS*:</b> .....	<b>91.6%</b>
<b>TOTAL:</b> .....	<b>100.0%</b>

Contains 0.66 lb of esfenvalerate per gallon.

\*CONTAINS XYLENE RANGE AROMATIC SOLVENT

EPA Reg. No. 352-515-66222

**KEEP OUT OF REACH OF CHILDREN  
WARNING-AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

**FIRST AID**

<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Do not give any liquid to the person.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

**NOTE TO PHYSICIAN:** If on skin, after drying apply vitamin E cream or oil if available. If not available, apply vegetable oil liberally over painful areas. The oil or cream may be used repeatedly until relief is achieved. Contains xylene range aromatic solvent – vomiting may cause aspiration pneumonia.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact PROSAR at 1-877-250-9291 24 hours a day.

For additional precautionary, handling and use statements, see inside of this booklet.



**M A N A** Raleigh, NC 27609

**Manufactured For:**  
**Makhteshim Agan**  
**of North America, Inc.**  
4515 Falls of Neuse Road  
Suite 300

EPA010505  
Adjourn071706

**Specimen Label**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
WARNING!**

May be fatal if swallowed. Harmful if inhaled. Do not get in eyes, on skin, or on clothing. Avoid breathing vapors or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

**PERSONAL PROTECTIVE EQUIPMENT**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical-resistance category selection chart.

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves, such as Barrier Laminate or Neoprene Rubber or Nitrile Rubber or Viton.
- Shoes plus socks.
- Protective eyewear.

Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

**USERS SHOULD:** Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**ENGINEERING CONTROL STATEMENTS**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR part 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning equipment or when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops if bees are visiting the treatment area.

**WILDLIFE PROTECTION IS EVERYONE'S RESPONSIBILITY**

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with the labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval.

The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Chemical-resistant gloves, such as Barrier Laminate or Neoprene Rubber or Nitrile Rubber or Viton.
- Shoes plus socks.
- Protective eyewear.

Adjourn™ Insecticide should be used only in accordance with recommendations in this label or in separate MAKHTESHIM AGAN OF NORTH AMERICA, INC. recommendations available through local dealers. MAKHTESHIM AGAN OF NORTH AMERICA, INC. will not be responsible for losses or damages resulting from (i) the reformulation of Adjourn™ Insecticide into any other product, and (ii) the use of Adjourn™ Insecticide in any manner that is not specifically recommended by MAKHTESHIM AGAN OF NORTH AMERICA, INC. on the label. User assumes all risk from such non-recommended use.

**GENERAL INFORMATION**

Adjourn™ Insecticide contains 0.66 pounds of active ingredient per gallon. For the applications given below, mix the required amount of Adjourn™ Insecticide in sufficient diluent to provide uniform coverage (Refer to Use Tables). Adjourn™ Insecticide may be applied by ground or aerial application equipment. For aerial application use the following directions unless otherwise specified in this label: use a minimum of 2 gallons per acre (gpa) of water, except in tree and orchard crops use a minimum of 10 gpa.

Apply at the recommended rates when insect populations reach locally determined economic thresholds. Consult the cooperative extension service, professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area.

**RESISTANCE**

Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

**INTEGRATED PEST MANAGEMENT**

MAKHTESHIM AGAN OF NORTH AMERICA, INC. recommends the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an Integrated Pest Management (IPM) program which can include biological, cultural, and genetic practices aimed at preventing economic pest damage. Application of this product should be based on IPM principles and practices including field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop or site systems in your area.

**SPRAY PREPARATION**

Spray equipment must be clean and free of previous pesticide deposits before applying Adjourn™ Insecticide.

**TANK MIXING AND COMPATIBILITY**

Do not tank mix Adjourn™ Insecticide with fungicides containing fentin hydroxide (triphenyltin hydroxide) such as "Super Tin" as crop injury may result. This product can be mixed with pesticide products labeled for use on crops on this label in accordance with the most restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing. Since formulations may be changed and new ones introduced, it is recommended that users premix a small quantity of a desired tank mix and observe for possible adverse changes (settling out, flocculation, etc.). Avoid mixtures of several materials and very concentrated spray mixtures. For best results, use of spray equipment having continuous agitation is recommended.

Adjourn™ Insecticide may be tank mixed with MAKHTESHIM AGAN OF NORTH AMERICA, INC. herbicide products when insect populations require control concurrent with the need for weed control. Follow all herbicide and Adjourn™ Insecticide label directions regarding proper usage. Adjourn™ Insecticide may be used in combination with 2,4-D herbicides providing that the following mixing directions are followed: 1) Do not apply the combination in a volume of water less than 2 gallons per acre total spray. 2) Always mix the Adjourn™ Insecticide thoroughly in the total volume of spray water first, followed by the addition of the 2,4-D herbicide. Because of the availability of a great variety of 2,4-D herbicide products, a test for physical compatibility should be conducted before field mixtures of a particular combination are made.

## CHEMIGATION

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) row, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system. Adjourntm Insecticide may be premixed in a supply tank with water, oil, fertilizer, or other appropriate tank mixed agricultural chemicals. A pretest of physical compatibility for untried tank mixes is advised. Agitation may be necessary. Application should be in sufficient water and of sufficient duration to apply the recommended rate evenly to the entire treated area. No run-off can be permitted during chemigation. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. Do not apply when windspeed favors drift beyond the area intended for treatment. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Do not connect an irrigation system (including greenhouse systems) used for Adjourntm Insecticide application to a public water system. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut-off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

## CROP ROTATION

ALL ROTATION CROPS MAY BE PLANTED IMMEDIATELY FOLLOWING LAST APPLICATION.

### SPRAY RECOMMENDATIONS AND PRECAUTIONS

- All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMANENT STREAMS, MARSHES, OR NATURAL PONDS; ESTUARIES AND COMMERCIAL FISH FARM PONDS.

- Do not apply by ground within 25 feet, or by air within 150 feet of lakes; reservoirs; rivers; permanent streams, marshes, or natural ponds; estuaries and commercial fish farm ponds. Increase the buffer zone to 450 feet when ultra low volume (ULV) application is made.
- For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of wing span or rotor diameter.
- Use the largest droplet size consistent with good pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orientating nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.
- Spray should be released at the lowest height consistent with pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.
- Make aerial or ground applications when the wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.
- Risk of exposure to aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.
- Do not cultivate within 10 feet of the aquatic area so as to allow growth of a vegetative filter strip.
- Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.
- Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

### SPRAY TANK CLEANOUT

Immediately following application of Adjourntm Insecticide, thoroughly clean all mixing and spray equipment. Flush the tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately). Take all necessary precautions when cleaning equipment. Do not clean near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

### SPECIFIC USES – FIELD CROPS

Crop	Insect	Application Rate		Acres treated per gal of Adjourntm	Last Application (days to harvest)
		lb ai/acre	fl oz/acre		
Corn (field)†	<b>Western Bean Cutworm</b>	0.015 – 0.03	2.9 – 5.8	44 – 22	21
	<b>Armyworm (True Armyworm)</b> <b>Black Cutworm (except CA)</b> <b>Chinch Bug</b> <b>Corn Earworm</b> <b>Corn Leaf Aphid</b> <b>Corn Rootworm (adult control)</b> <b>Cutworm</b> <b>Flea Beetle</b> <b>Grasshopper</b> <b>Oat Bird-Cherry Aphid</b> <b>Southwestern Corn Borer</b> <b>Stalk Borer</b>	0.03 – 0.05	5.8 – 9.6	22 -13	
	<b>European Corn Borer</b>	0.04 – 0.05	7.8 – 9.6	16 – 13	
	<p><b>Black Cutworm</b> – Adjourntm may be applied at 3.2 – 9.6 fl oz/acre (0.0165 – 0.05 lb ai/acre) for the control of black cutworm when applied at planting of corn (except CA).</p> <p><b>Chinch Bug</b> – Over the top sprays, as used for control of ear and stalk infesting corn pests, are not adequate for chinch bug control. It is very important that the spray be directed at the base of the plant through the use of drop nozzles or some other mechanism.</p> <p><b>Corn Earworm</b> – First application should be at or before silking. Repeat applications may be applied if economically damaging populations exist. Subsequent applications should be made at 3-5 day intervals until silking is completed.</p> <p><b>Corn Leaf Aphid, Oat Bird-Cherry Aphid</b> – For optimum results, direct the spray at the aphid population so as to achieve maximum coverage of the exposed insects. Aphids not contacted by the spray, such as in whorls and leaf axils, may not be adequately controlled.</p> <p><b>Corn Rootworm (Adult)</b> – Apply at the first sign of silk feeding. Adjourntm may be tank-mixed with methyl parathion and applied on field corn where supplemental control of Adult Corn Rootworm is desired in conjunction with insects controlled by Adjourntm when used alone. Refer to the Adjourntm and methyl parathion labels for appropriate rates of the individual products for controlling the respective insects.</p> <p><b>Cutworm</b> – Applications for cutworm control may be applied before, during, or after planting as required to protect emerging or emerged corn seedlings.</p> <p><b>European Corn Borer</b> <b>First brood:</b> Spray while eggs are in the blackhead stage or before the larvae enter the whorl. Application by ground equipment is suggested. Good coverage of both upper and lower leaf surfaces is essential. This can be accomplished with drop nozzles over the row and on each side of the corn plant. Multiple applications may be required when egg laying is prolonged or where moderate to heavy populations are present. A higher rate is recommended for moderate to heavy populations. Proper coverage by ground equipment usually requires 20-30 gallons of carrier. Once larvae enter the whorl, foliar sprays will not provide adequate control.</p>				

Crop	Insect	Application Rate		Acres treated per gal of Adjourn™	Last Application (days to harvest)
		lb ai/acre	fl oz/acre		
<b>Corn (field)†</b> <i>(cont'd)</i>	<p><b>Second brood:</b> Make applications when sufficient egg masses are found. Spray when eggs are in the blackhead stage or starting to hatch. When egg laying is prolonged or a third generation is present, additional sprays may be required. A higher rate is recommended for moderate to heavy populations. Good coverage above, below, and in the ear zone is essential. This usually requires 2 – 3 gallons of carrier by air. If ground equipment is used, drop nozzles on each side of the plant will provide best coverage. Adjourn™ may be tank-mixed with methyl parathion and applied on field corn where supplemental control of European Corn Borer is desired in conjunction with insects controlled by Adjourn™ when used alone.</p> <p>Refer to the Adjourn™ and methyl parathion labels for appropriate rates of the individual products for controlling the respective insects.</p> <p><b>Southwestern Corn Borer</b> – For moderate to heavy infestations, higher rates (0.036-0.05 lb ai per acre) are recommended.</p> <p><b>Stalk Borer, Flea Beetle</b> – Application must be made early in migration from grassy areas to corn, before borers enter the plant.</p> <p><b>Western Bean Cutworm</b> – Apply before larvae enter the ear.</p> <p>† Do not apply more than 0.25 lb ai per acre per season.</p>			22	21
<b>Corn (pop)</b>	<p>For specific insect control recommendations refer to Field Corn (above). Follow directions carefully.</p> <p>Multiple applications and/or shortened intervals between sprays must be used to insure proper insect control. Do not apply more than 0.5 lb a.i. per acre per season.</p>				1
<b>Corn (seed)</b>	<p>For specific insect control recommendations refer to Field Corn (above). Follow directions carefully.</p> <p>Multiple applications and/or shortened intervals between sprays must be used to insure proper insect control. Do not apply more than 0.25 lb a.i. per acre per season.</p>				1
<b>Cotton</b>	<b>Cotton Leaf Perforator</b>	0.03	5.8	22	21
	<b>Beet Armyworm*</b> <b>Black Cutworm</b> (except CA) <b>Boll Weevil</b> <b>Cabbage Looper</b> <b>Cotton Aphid*</b> <b>Cotton Bollworm</b> <b>Cotton Leafworm</b> <b>Cutworms</b> <b>Fleahoppers</b> <b>Grasshoppers</b> <b>Green Stink Bug</b> (except CA) <b>Leafhoppers</b> <b>Lygus Bugs</b> <b>Pink Bollworm</b> <b>Plant Bugs</b> <b>Saltmarsh Caterpillar</b> <b>Southern Green Stink Bug</b> (except CA) <b>Thrips</b> (on seedling cotton) <b>Tobacco Budworm</b> <b>Whitefly*</b>	0.03 – 0.05	5.8 – 9.6	22 – 13	
	<b>NOTE:</b> For light infestations of the above insects	0.02	3.9	33	
	<p>* Aids in control. May be applied in water or non-volatile vegetable oils. When applying Adjourn™ in an oil carrier, apply a total spray volume of at least 1 qt per acre. When applying Adjourn™ in a water carrier, apply at least 1 gal per acre by air (at least 3 gal per acre in Arizona and 5 gal per acre in California) or 4 gal per acre by ground. Do not apply more than 0.5 lb ai per acre per season. Do not graze livestock on treated fields or feed treated trash. <b>Black Cutworm</b> – Adjourn™ may be applied at 3.2 – 9.6 fl oz/acre (0.0165 – 0.05 lb ai/acre) for the control of black cutworm when applied at planting of cotton (except CA). <b>Boll Weevil</b> – To control Boll Weevil infestations, a 3- to 5-day interval between applications may be necessary. <b>Heliopsis spp.</b> – Adjourn™ can provide contact ovicidal effect on Heliopsis spp. eggs when applied according to label directions for control of tobacco budworm; application should be timed to correspond with peak egg deposition to achieve maximum ovicidal effect. Use on this pest stage (egg) is not registered in California. Do not make more than a total of 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season. Synthetic pyrethroid products include Adjourn™ insecticide, "Ambush" insecticide, "Ammo" insecticide, "Asana XL" insecticide, "Baythroid" emulsifiable pyrethroid insecticide, "Capture" insecticide/miticide, "Danitol 2.4 EC" spray insecticide/miticide, "Decis" insecticide, "Fury" insecticide, "Karate" insecticide, "Mustang" insecticide, "Pounce" insecticide and "Scout X-tra" insecticide.</p>				
<b>Peanuts</b>	<b>Corn Earworm</b> <b>Potato Leafhopper</b> <b>Red-necked Peanut Worm</b> <b>Velvetbean Caterpillar</b>	0.015 – 0.03	2.9 – 5.8	44 – 22	21
	<b>Beet Armyworm*</b> <b>Cutworms</b> <b>Granulate Cutworm</b> <b>Grasshoppers</b>	0.03 – 0.05	5.8 – 9.6	22 – 13	
	<b>Fall Armyworm*</b> <b>Lesser Cornstalk Borer**</b>	0.05	9.6	13	
	<p>* Aids in control. For heavy infestations use of DuPont™ LANNATE® insecticide is recommended. Please refer to the LANNATE® label for use instructions. ** Aids in control. Do not feed or graze livestock on treated vines. Do not apply more than 0.15 lb a.i. per acre per season.</p>				

Crop	Insect	Application Rate		Acres treated per gal of Adjourn™	Last Application (days to harvest)
		lb ai/acre	fl oz/acre		
Sorghum (Grain) (except CA)	Sorghum Midge	0.015 – 0.03	2.9 – 5.8	44 – 22	21
	Black Cutworm Chinch Bugs Corn Earworm (headworm) Cutworms	0.03 – 0.05	5.8 – 9.6	22 – 13	
	Do not apply more than 0.15 lb a.i. per acre per season. When applying in non-volatile vegetable oils use a total spray volume of 1 or more qts per acre. <b>Black Cutworm</b> – Adjourn™ may be applied at 3.2 – 9.6 fl oz/acre (0.0165 – 0.05 lb ai/acre) for the control of black cutworm when applied at planting of sorghum. <b>Chinch Bug Control</b> – For optimum results, spray should be directed at base of plants.				
Soybean	Green Cloverworm Mexican Bean Beetle Potato Leafhopper Saltmarsh Caterpillar Velvetbean Caterpillar Woollybear Caterpillar	0.015 – 0.03	2.9 – 5.8	44 – 22	21
	Bean Leaf Beetle Beet Armyworm* Cabbage Looper Corn Earworm Cutworms Grasshoppers Green Stink Bug (except CA) Japanese Beetle (adult) Southern Green Stink Bug Soybean Aphid (except CA) Three-cornered Alfalfa Hopper	0.03 – 0.05	5.8 – 9.6	22 – 13	
	* Aids in control. When applying in non-volatile vegetable oils, use a total spray volume of at least 1 qt. Do not feed or graze livestock on treated fields. Do not apply more than 0.2 lb ai per acre per season.				
Sugar Beets	Beet Armyworm* Beet Webworm Cabbage Looper Cutworms (seedling spray) Flea Beetle (except CA) Grasshoppers Leafhoppers Saltmarsh Caterpillar	0.03 – 0.05	5.8 – 9.6	22 – 13	21
	* Aids in control. Do not apply more than 0.15 lb ai/acre per season. Apply with ground or air equipment using sufficient water to provide uniform coverage (minimum of 2 gal of water per acre).				
Sugarcane	Sugarcane Borer	0.03 – 0.05	5.8 – 9.6	22 – 13	21
	Do not apply more than 0.2 lb ai/acre per season.				
Sunflower	Sunflower Beetle (except CA)	0.0075 – 0.03	1.45 – 5.8	88 – 22	28
	Banded Sunflower Moth Beet Armyworm* Cutworms Grasshoppers Heliothis (complex) Leafhoppers Sunflower Maggot Sunflower Moth Sunflower Seed Weevil Sunflower Stem Weevil	0.03 – 0.05	5.8 – 9.6	22 – 13	
	* Aids in control. Do not apply more than 0.2 lb. ai/acre per season.				

**SPECIFIC USES – FRUITS**

**DILUTE SPRAY:** Apply specified dosage per 100 gallons of water in a uniform spray applied to the point of drip with conventional ground equipment. Do not exceed maximum number of gallons per acre indicated.

**NOTE:** In order to apply the correct amount of Adjourn™ Insecticide to your orchard you must know the number of gallons of water needed to spray one acre of your trees to the point of drip. If you do not already know this gallonage, you should conduct a test to determine it. If you do not know how to conduct such a test with your equipment, you should ask for assistance from your equipment dealer or State Extension specialist.

**CONCENTRATE SPRAY:** APPLY SPECIFIED DOSAGE PER ACRE IN NO LESS THAN 30 GALS. OF WATER PER ACRE BY GROUND. FOR AERIAL APPLICATION IN TREE AND ORCHARD CROPS, USE A MINIMUM OF 10 GALLONS OF WATER PER ACRE. WHEN APPLYING ADJOURN™ BY AIR, CONSULT YOUR COOPERATIVE EXTENSION SERVICE FOR APPLICATION GUIDELINES.

Crop	Insect	Application Rate			Acres treated per gal of Adjourn™	Last Application (days to harvest)
		Concentrate Spray		Dilute Spray		
		lb ai/acre	fl oz/acre	fl oz/100 gal		
Apples	<b>Apple Aphid</b> <b>Apple Maggot</b> <b>Codling Moth</b> <b>Green Fruitworm</b> <b>Lesser Appleworm</b> <b>Mullein Plant Bug</b> (except CA) <b>Oblique Banded Leafroller</b> <b>Oriental Fruit Moth</b> <b>Periodical Cicada</b> <b>Plant Bugs</b> (Tarnished Plant Bug, Stink Bugs) <b>Plum Curculio</b> <b>Red – Banded Leafroller</b> <b>Rosy Apple Aphid</b> <b>San Jose Scale</b> (fruit infestations only) <b>Tentiform Leaf Miner</b> <b>Tufted Apple Bud Moth</b> <b>Variigated Leafroller</b> <b>White Apple Leafhopper</b>	0.025 – 0.075	4.8 – 14.5	2.0 – 5.8	26 – 9	21
	<b>Apple Ermine Moth</b> (ID, OR & WA only)			3.0		
	<b>Tufted Apple Bud Moth (overwintering)</b> (MD, NC, NJ, PA, VA, WV only)	0.04 – 0.075	8.0 – 14.5		16 – 9	
	Do not feed or graze livestock on treated orchard floors. Do not apply more than 0.525 lbs. a.i. per acre per season. For dilute spray apply 200 – 600 gals per acre, but do not apply more than 14.5 fl. oz. of Adjourn™ per acre per treatment. <b>Apple Ermine Moth</b> – Apply with 2 to 4 gallons of superior spray oil in 100 gallons of water in a spray-to-wet application to insure thorough coverage of all stems and branches where Apple Ermine Moth hibernacula are found. When using on apple nursery stock, do not treat bundled plants since it is difficult to achieve a full coverage application which could result in less than complete control. Make first application in the fall after 90% of leaf fall has occurred – usually after October 15. Make a second application 7 to 14 days later. <b>NOTE:</b> Overwintering larvae do not die until approximately 30 days after application. <b>Plant Bug, Rosy Apple Aphid Control</b> – Time of application is critical in achieving control. Use prebloom and post bloom spray timings recommended by State Extension Services. <b>Tufted Apple Bud Moth (overwintering)</b> – For use on apple for the control of overwintering larvae of the tufted apple bud moth with directed ground application to the apple orchard floor. Make one application of Adjourn™ at either pink stage of apple or at petal fall stage of apple. Use the lower rate on small larvae (pink stage of apple) and/or on lower populations. Use the higher rate on larger larvae (petal fall stage of apple) and/or on moderate to high populations. Apply specified dosage per acre to the orchard floor in no less than 30 gals of water per acre by ground to obtain uniform coverage. Apply treatment in a band from trunk to drip line to allow coverage of areas where overwintering tufted apple bud moth are found. <b>Beneficial Insects:</b> Application of Adjourn™ to the groundcover at the pink stage of apple development may be toxic to overwintering <i>Stethorus punctum</i> . <i>S. punctum</i> is a coccinellid insect and the major predator of spider mites in the MD, NC, NJ, PA, VA and WV fruit growing areas. This predator overwinters in the same areas of the orchard groundcover as the tufted apple bud moth and moves into apple trees from mid-April through mid-May when maximum daily temperatures exceed 68°F. Emergence from the groundcover is 20-70% complete by the pink stage and 90-100% complete by petal fall on the apple cultivar Yorking.					
Blueberry (except CA)	<b>Aphids</b> (NJ only) <b>Blueberry Spanworm</b> <b>Cherry Fruitworm*</b> <b>Cranberry Fruitworm</b> <b>Cranberry Weevil*</b> <b>Grasshoppers</b> <b>Japanese Beetle</b> <b>Leafhoppers</b> <b>Red Striped Fireworm*</b>	0.025 – 0.05	4.8 – 9.6		26 – 13	14
	<b>Blueberry Maggot</b> <b>Black Vine Weevil</b> (adult control) <b>Strawberry Root Weevil</b> (adult control) (OR, WA only)	0.05	9.6		13	
	* Aids in control. Do not apply more than 0.2 lb ai/acre per season (38.4 fluid ounces of product per acre per season). Use of ground application is recommended; for ground application use a minimum of 50 gals of water per acre. Do not apply this product through any type of irrigation system. <b>NOTE:</b> Adjourn™ can act as a bee repellent, do not apply within 7 days of pollination. Apply as a pre-bloom or post-bloom spray only.					

Crop	Insect	Application Rate			Acres treated per gal of Adjourn™	Last Application (days to harvest)
		Concentrate Spray		Dilute Spray		
		lb ai/acre	fl oz/acre	fl oz/100 gal		
<b>Blueberry</b> (except CA) (cont'd)	<b>Black vine weevil &amp; strawberry root weevil (adult control) (OR, WA only):</b> Look for leaf notching beginning in late May to early June as the first sign of weevil feeding. Also check for adults on or just below the soil surface around the base of plants. Apply Adjourn™ within two to three weeks of first sign of infestation. Do not apply by air; apply by ground using a minimum of 50 gallons of water per acre. Direct spray to provide full coverage of foliage and soil area around base of plants. Best results are from applications made after dark when temperatures are warm and weevils are actively feeding. Root weevils emerge over a several week period, make additional applications when signs of new feeding appear.				14	
<b>Caneberries</b> (blackberries, boysenberries, dewberries, loganberries, raspberries, youngberries, and varieties of these) (except CA)	<b>Aphids</b> <b>Oblique Banded Leafroller</b> <b>Orange Tortrix</b>	0.025 – 0.05	4.8 – 9.6		26 – 13	7
	<b>Adult Root Weevils*</b> (OR & WA only)	0.05	9.6		13	
<p>*Aids in control. Do not apply by air. Do not apply more than 0.15 lb ai per acre per season. Do not apply this product through any type of irrigation system. <b>NOTE:</b> Adjourn™ can act as a bee repellent, do not apply within 7 days of pollination. Apply as a pre-bloom or post-bloom spray only. Remove bees prior to application. For maximum safety to bees, apply Adjourn™ in the evening after sunset. <b>Adult Root Weevils (OR &amp; WA only):</b> Look for leaf notching beginning in late May to early June as the first sign of weevil feeding. Also check for adults on or just below the soil surface around the base of plants. Apply Adjourn™ within two to three weeks of infestation. Apply by ground using a minimum of 50 gallons of water per acre. Direct spray to provide thorough coverage of all plant foliage and the soil area around the base of the plant. Best results are from applications made after dark when temperatures are warm and weevils are actively feeding. Root weevils emerge over a period of several weeks; make subsequent applications when signs of new feeding appear. <b>Oblique Banded Leafroller Orange Tortrix and Aphids:</b> Apply as a full coverage spray in a minimum of 50 gallons of water with ground equipment only. Apply no earlier than 12 days before harvest and no later than 7 days before harvest.</p>						
<b>Kiwifruit</b>	<b>Boxelder Bug</b> (suppression only)	0.05	9.6	–	13	14
	Spray in sufficient water for thorough coverage. A maximum of 7 applications is allowed per season (total of 0.35 lb ai/acre/season) with a minimum of 7 days between treatments.					
<b>Pear</b>	<b>Codling Moth</b> <b>Green Fruitworm</b> <b>Leafrollers</b> <b>Pear Psylla</b> <b>Pear Slug</b> <b>Periodical Cicada</b> <b>Plum Curculio</b>	0.025 – 0.075	4.8 – 14.5	2.0 – 5.8	26 – 9	28
	Do not apply more than 0.375 lbs. a.i. per acre per season. Do not apply more than 0.225 lb. a.i. per acre between bloom and harvest. Do not feed or graze livestock on treated orchard floors. For dilute spray apply 200 – 600 gals per acre, but do not apply more than 14.5 fl. oz. of Adjourn™ per acre per treatment.					
<b>Pear</b> (Dormant)	<b>Pear Psylla</b>	0.05 – 0.1	9.6 – 19.2	7.3 – 12.8	13.2 – 6.6	
	Apply during dormant to prebloom (white bud) stage only. Do not apply more than 0.2 lb a.i. per acre per season. Do not graze orchard floor. For dilute spray apply 150 – 250 gals per acre, but do not apply more than 19.2 fl. oz. of Adjourn™ per acre per treatment.					
<b>Stone Fruits</b> (including apricots, cherries, nectarines, peaches, plums, prune plums)	<b>American Plum Borer</b> <b>Black Cherry Aphid</b> <b>Cherry Fruit Fly</b> <b>Green Fruitworm</b> <b>Leafhoppers</b> <b>Leafrollers</b> <b>Lesser Peach Tree Borer</b> <b>Oriental Fruit Moth</b> <b>Peach Tree Borer</b> <b>Peach Twig Borer</b> <b>Periodical Cicada</b> <b>Plant Bugs</b> (Tarnished Plant Bug, Stink Bugs) <b>Plum Curculio</b> <b>Western Cherry Fruit Fly</b>	0.025 – 0.075	4.8 – 14.5	2.0 – 5.8	26 – 9	14
	<b>Peach Twig Borer</b> (Dormant) (CA only)	0.04 – 0.075	8.0 – 14.5	3.1 – 5.8	16 – 9	
	<b>Tufted Apple Bud Moth</b> (overwintering) (Peach only) (MD, NC, NJ, PA, VA and WV only)	0.04 – 0.075	8.0 – 14.5		16 – 9	
	Do not apply more than 0.375 lbs a.i. per acre per season with no more than 0.3 lbs. a.i. per acre per season between bloom and harvest. For dilute spray apply 200 – 400 gals per acre, but do not apply more than 14.5 fl. oz. of Adjourn™ per acre per treatment. Do not graze livestock on treated orchard floors. <b>Peach Tree Borer, American Plum Borer, Lesser Peach Tree Borer Control</b> – Apply as directed trunk and scaffold limb spray. Thorough coverage of trunk and scaffold limbs is required. <b>Peach Twig Borer (Dormant)</b> – Make application with an EPA registered dormant oil; for specific recommendations on use of oil consult manufacturer's label. For best performance, ground application equipment is recommended. <b>Peach Twig Borer, Plum Curculio, Oriental Fruit Moth, Cherry Fruit Fly, Western Cherry Fruit Fly, Leafrollers, Black Cherry Aphid, Periodical Cicada Control</b> – Apply by ground sprayer to achieve thorough coverage of all aerial portions of the tree. <b>Plant Bug Control</b> – Time of application is critical in achieving control. Use pre-bloom and postbloom spray timings recommended by State Extension Services.					

Crop	Insect	Application Rate			Acres treated per gal of Adjourn™	Last Application (days to harvest)
		Concentrate Spray		Dilute Spray		
		lb ai/acre	fl oz/acre	fl oz/100 gal		
Stone Fruits (cont'd)	<p><b>Tufted Apple Bud Moth (overwintering)</b> – For use on peach for the control of overwintering larvae of the tufted apple bud moth with directed ground application to the peach orchard floor. Make one application of Adjourn™ at either popcorn stage of peach or at shucksplit stage of peach.</p> <p>Use the lower rate on small larvae (popcorn stage of peach) and/or on lower populations. Use the higher rate on larger larvae (shucksplit stage of peach) and/or on moderate to high populations.</p> <p>Apply specified dosage per acre to the orchard floor in no less than 30 gals of water per acre by ground to obtain uniform coverage. Apply treatment in a band from trunk to drip line to allow coverage of areas where overwintering tufted apple bud moth are found.</p> <p><b>Beneficial Insects:</b> Application of Adjourn™ to the groundcover at the popcorn stage of peach development may be toxic to overwintering <i>Stethorus punctum</i>. <i>S. punctum</i> is a coccinellid insect and the major predator of spider mites in the MD, NC, NJ, PA, VA and WV fruit growing areas. This predator overwinters in the same areas of the orchard groundcover as the tufted apple bud moth and moves into peach trees from mid-April through mid-May when maximum daily temperatures exceed 68°F. Emergence from the groundcover is 90-100% complete by shucksplit. Shucksplit is the preferred timing to minimize predator toxicity.</p>				14	

**SPECIFIC USES – TREE NUT CROPS**

**DILUTE SPRAY:** Apply specified dosage per 100 gallons of water in a uniform spray applied to the point of drip with conventional ground equipment. Do not exceed maximum number of gallons per acre indicated.

**NOTE:** In order to apply the correct amount of Adjourn™ Insecticide to your orchard you must know the number of gallons of water needed to spray one acre of your trees to the point of drip. If you do not already know this gallonage, you should conduct a test to determine it. If you do not know how to conduct such a test with your equipment, you should ask for assistance from your equipment dealer or State Extension specialist.

**CONCENTRATE SPRAY:** APPLY SPECIFIED DOSAGE PER ACRE IN NO LESS THAN 30 GALS. OF WATER PER ACRE BY GROUND. FOR AERIAL APPLICATION IN TREE AND ORCHARD CROPS, USE A MINIMUM OF 10 GALLONS OF WATER PER ACRE. WHEN APPLYING ADJOURN™ BY AIR, CONSULT YOUR COOPERATIVE EXTENSION SERVICE FOR FURTHER APPLICATION GUIDELINES.

Crop	Insect	Application Rate			Acres treated per gal of Adjourn™	Last Application (days to harvest)
		Concentrate Spray		Dilute Spray		
		lb ai/acre	fl oz/acre	fl oz/100 gal		
Almonds	<p><b>Navel Orangeworm</b> <b>Peach Twig Borer</b> <b>Peach Twig Borer (Dormant)</b> (CA only)</p>	0.05 – 0.1	9.6 – 19.2	7.3 – 12.8	13.2 – 6.6	21
	<p>Do not apply more than 0.2 lb. a.i. per acre per season. Do not graze livestock on treated orchard floors. <b>Peach Twig Borer (Dormant)</b> – Make application with an EPA registered dormant oil; for specific recommendations on use of oil consult manufacturer's label. For best performance, ground application equipment is recommended.</p>					
Filberts	<p><b>Filbertworm</b> <b>Oblique Banded Leafroller</b></p>	0.05 – 0.1	9.6 – 19.2	7.3 – 12.8	13.2 – 6.6	21
	<p>Make first application after emergence of filbert worm moths in early summer. Do not apply a second treatment earlier than three weeks after the first. Do not apply more than 0.2 lb. a.i. per acre per season. For dilute spray, apply 200 – 400 gals. per acre, but do not apply more than 19.2 fl. oz. of Adjourn™ per acre per treatment. Do not graze livestock on treated orchard floors.</p>					
Pecans	<p><b>Hickory Shuckworm</b> <b>Pecan Aphids</b> <b>Pecan Nut Casebearer</b> <b>Pecan Leaf Phylloxera</b> <b>Pecan Spittlebug</b> <b>Pecan Stem Phylloxera</b> <b>Pecan Weevil</b></p>	0.025 – 0.075	4.8 – 14.5	2.0 – 5.8	26 – 9	21
	<p>Do not feed or graze livestock on treated orchard floors. Do not apply more than 0.3 lbs. a.i. per acre per season. For dilute spray apply 200 – 600 gals per acre, but do not apply more than 14.5 fl. oz. of Adjourn™ per acre per treatment. <b>Phylloxera</b> – Correct timing of spray applications is critical in achieving optimum control of leaf and stem phylloxera. Consult local spray recommendations for correct times of application.</p>					
Walnuts	<p><b>Codling Moth</b> <b>Navel Orangeworm</b> <b>Walnut Aphid</b> <b>Walnut Husk Fly</b></p>	0.05 – 0.1	9.6 – 19.2	4.0	13 – 6	21
	<p>Do not apply more than 0.2 lb. a.i. per acre per season. For dilute sprays, apply 200 – 400 gals per acre. Do not feed or graze livestock on treated crop floors. <b>NOTE:</b> Use of baits in Walnut Husk fly sprays is recommended where endorsed by local Agricultural Extension Service.</p>					

**SPECIFIC USES – VEGETABLE CROPS**

Crop	Insect	Application Rate		Acres treated per gal of Adjour™	Last Application (days to harvest)
		lb ai/acre	fl oz/acre		
Artichoke	<b>Artichoke Plume Moth</b>	0.03 – 0.05	5.8 – 9.6	22 – 13	1
	Do not apply more often than each 14 days. Apply no more than 0.15 lb. ai/acre between bud formation and harvest of an individual fruit. Apply in a minimum of 10 gallons per acre by air and 50 – 200 gallons per acre by ground (use sufficient water to obtain coverage without excessive runoff).				
Beans, Dry (Including adzuki bean, blackeyed pea, broad bean (dry), chickpea, cow pea, crowder pea, field bean, kidney bean, lima bean (dry), mung bean, navy bean, pinto bean, southern pea, tepary bean) Peas, Dry Lentils	<b>Leafhoppers (except CA)</b> <b>Mexican Bean Beetle</b> <b>Saltmarsh Caterpillar</b>	0.015 – 0.03	2.9 – 5.8	44 – 22	21
	<b>Beet Armyworm*</b> <b>Cabbage Looper</b> <b>Corn Earworm</b> <b>Corn Rootworm (Adults)</b> <b>Cowpea Curculio</b> <b>Cucumber Beetle</b> <b>Cutworms</b> <b>Grasshoppers</b> <b>Green Cloverworm</b> <b>Leafhoppers</b> <b>Painted Lady Butterfly (larvae)</b> <b>Pea Aphid</b> <b>Potato Leafhopper</b> <b>Soybean Aphid (except CA)</b> <b>Velvetbean Caterpillar</b> <b>Western Bean Cutworm</b>	0.03 – 0.05	5.8 – 9.6	22 – 13	
	<b>Pea Leaf Weevil (ID, OR &amp; WA only)</b> <b>Pea Weevil (ID, OR &amp; WA only)</b>	0.025 – 0.05	4.8 – 9.6	26 – 13	
	*Aids in control. Do not apply more than 0.2 lb a.i. per acre per season. Do not feed or graze livestock on treated vines. <b>Pea Weevil &amp; Pea Leaf Weevil (ID, OR &amp; WA)</b> – Time of application is critical in achieving control of pea weevil. For optimum results, apply at bloom prior to detecting adult pea weevils. Once adult pea weevil populations reach a level of 2 or more adults per 25 sweeps, control may be reduced.				
Beans, Snap Also known as: (blue lake, bush, common, edible-podded, filet, flageolet, French, French horticultural, frijoles comunes, garden, green, haricot, haricot commun, Italian, judia comum, Kentucky wonder, magic, pole, romano, string, succulent, vainica, wax)	<b>Leafhoppers (except CA)</b> <b>Mexican Bean Beetle</b> <b>Saltmarsh Caterpillar</b>	0.015 – 0.03	2.9 – 5.8	44 – 22	3
	<b>Beet Armyworm*</b> <b>Cabbage Looper</b> <b>Corn Earworm</b> <b>Corn Rootworm (adults)</b> <b>Cucumber Beetle</b> <b>Cucumber Beetle (adults)</b> <b>Cutworm (seedling spray)</b> <b>European Corn Borer</b> <b>Flea Beetle</b> <b>Grasshoppers</b> <b>Green Cloverworm</b> <b>Leafhopper</b> <b>Leafminer (Guam only)</b> <b>Pea Aphid</b> <b>Potato Leafhopper</b> <b>Soybean Aphid (except CA)</b> <b>Velvet Bean Caterpillar</b> <b>Western Bean Cutworm</b>	0.03 – 0.05	5.8 – 9.6	22 – 13	
*Aids in control. Do not apply more than 0.2 lb ai per acre per season. Do not allow livestock to graze treated bean fields. Do not harvest treated bean vines for livestock forage, fodder, or hay.					
Broccoli, (including chinese broccoli) Cabbage, Cauliflower, Chinese Cabbage (tight headed varieties only)	<b>Imported Cabbageworm</b>	0.015 – 0.03	2.9 – 5.8	44 – 22	3
	<b>Alfalfa Looper</b> <b>Beet Armyworm*</b> <b>Cabbage Looper</b> <b>Cutworm</b> <b>Flea Beetle</b> <b>Grasshoppers</b>	0.03 – 0.05	5.8 – 9.6	22 – 13	
* Aids in control. Do not apply more than 0.4 lbs. a.i. per acre per season.					
Carrots	<b>Aster Leafhopper</b> <b>Cutworms</b> <b>Leafhoppers</b>	0.03 – 0.05	5.8 – 9.6	22 – 13	7
	<b>Carrot Weevil</b>	0.05	9.6	13	
Do not apply more than 0.5 lb ai/acre per season. For aerial application apply in a minimum of 5 gals of water per acre. Thorough spray coverage of crown area is essential. Use of ground application is recommended. <b>Carrot Weevil</b> – Begin treatment when weevils become active.					

Crop	Insect	Application Rate		Acres treated per gal of Adjournal™	Last Application (days to harvest)
		lb ai/acre	fl oz/acre		
Collards	Alfalfa Looper Beet Armyworm* Cabbage Looper Cutworm Flea Beetle Grasshopper Imported Cabbageworm	0.03 – 0.05	5.8 – 9.6	22 – 13	7
	* Aids in control. Do not feed livestock on treated plant parts. Do not apply more than 0.2 lbs. a.i. per acre per season. For aerial application apply in a minimum of 5 gals of water per acre.				
Cucumber, Melons, (cantaloupe, honeydew melons, muskmelon, watermelon), Pumpkin, Squash (summer, winter)	Cabbage Looper Corn Earworm Cucumber Beetle (adults) Cutworms (seedling spray) Grasshoppers Leafhoppers Pickleworm Plant Bugs (Lygus Bugs, Stink Bugs) Rindworms Squash Bug Squash Vine Borer	0.03 – 0.05	5.8 – 9.6	22 – 13	3
	Do not apply more than 0.25 lbs. a.i. per acre per season.				
Eggplant	Colorado Potato Beetle Corn Earworm European Corn Borer Flea Beetles Loopers	0.03 – 0.05	5.8 – 9.6	22 – 13	7
	Apply when insects are observed or when insect damage is observed. Repeat applications at 7- to 10-day intervals to achieve control. Do not apply more than 0.35 lbs. a.i. per acre per season.				
Kohlrabi (TX only)	Cabbage Looper	0.03 – 0.05	5.8 – 9.6	22 – 13	3
	Do not apply more than 0.4 lbs. a.i. per acre per season.				
Lentils	See "Beans, Dry"				
Lettuce, Head AZ, CA, CO, FL, NM & TX ONLY	Alfalfa Looper Beet Armyworm* Cabbage Looper Heliothis spp.	0.025 – 0.05	4.8 – 9.6	26 – 13	7
	* Aids in control. Do not apply more than 0.35 lbs. a.i. per acre per season.				
Mustard Greens	Cabbage Looper Imported Cabbageworm	0.05	9.6	13	7
	Do not apply more than 0.2 lbs/a.i. per acre per season.				
Peas, Dry	See "Beans, Dry"				
Peas, Green	Green Cloverworm Pea Aphid	0.015 – 0.03	2.9 – 5.8	44 – 22	3
	Alfalfa Caterpillar Alfalfa Looper Armyworm Cabbage Looper Celery Looper Corn Earworm Cutworms Imported Cabbageworm	0.03 – 0.05	5.8 – 9.6	22 – 13	
	Pea Leaf Weevil (ID, OR & WA only) Pea Weevil (ID, OR & WA only)	0.025 – 0.05	4.8 – 9.6	26 – 13	
	Do not apply more than 0.1 lb ai/acre per season. Do not feed treated pea vines to livestock. Pea Weevil & Pea Leaf Weevil (ID, OR & WA only) – Time of application is critical in achieving control of pea weevil. For optimum results, apply at bloom prior to detecting adult pea weevils. Once adult pea weevil populations reach a level of 2 or more adults per 25 sweeps, control may be reduced.				
Pepper	Beet Armyworm* Colorado Potato Beetle Corn Earworm Cucumber Beetle (adults) European Corn Borer Flea Beetles Loopers Pepper Weevil* Southern Armyworm	0.03 – 0.05	5.8 – 9.6	22 – 13	7
	* Aids in control. Apply when insects are observed or when insect damage is observed. Repeat applications at 7- to 10-day intervals to achieve control. Do not apply more than 0.35 lbs. a.i. per acre per season.				

Crop	Insect	Application Rate		Acres treated per gal of Adjournal™	Last Application (days to harvest)	
		lb ai/acre	fl oz/acre			
Potato	Leafhoppers (except CA) Potato Psyllid Potato Tuberworm	0.015 – 0.03	2.9 – 5.8	44 – 22	7	
	Beet Armyworm* Buckthorn Aphid Cabbage Looper Colorado Potato Beetle Cucumber Beetle (adult control) Cutworms European Corn Borer Flea beetles Grasshoppers Potato Aphid Potato Leafhopper Tarnished Plant Bug Western Yellow-Striped Armyworm	0.03 – 0.05	5.8 – 9.6	22 – 13		
	Colorado Potato Beetle (Long Island, NY only)	0.05	9.6	13		
	Aids in control. Do not apply more than 0.35 lb. a.i. per acre per season.					
Radishes	Armyworms Beetles	0.03 – 0.05	5.8 – 9.6	22 – 13	7	
	Do not apply more than 0.1 lb. ai/acre per season.					
Sweet Corn†	Western Bean Cutworm	0.015 – 0.03	2.9 – 5.8	44 – 22	1	
	For additional information consult directions for use under field corn.					
	Armyworm Banded Cucumber Beetle Beet Armyworm* Chinch Bugs Corn Earworm Corn Leaf Aphid Corn Rootworms (adults) Cutworms European Corn Borer Flea Beetles Grasshoppers Oat Bird-Cherry Aphid Sap Beetles (adults) Southwestern Corn Borer Stalk Borer Tarnished Plant Bug	0.03 – 0.05	5.8 – 9.6	22 – 13		
	Corn Silkfly (except CA)**	0.05	9.6	13		
<p>* Aids in control. ** Suppression only. <b>For Ear Protection</b> – Begin applications either just before or at time of silking. For additional information consult directions for use under field corn. <b>Corn Leaf Aphid &amp; Oat Bird-Cherry Aphid</b> – For optimum results, direct the spray at the aphid population so as to achieve maximum coverage of the exposed insects. Aphids not contacted by the spray, such as in whorls and leaf axils, may not be adequately controlled. <b>Corn Silkfly (except CA)</b> – Direct application to the ear zone to obtain thorough coverage of the corn silk. † Do not apply more than 0.5 lb a.i. per acre per season.</p>						
Tomato	Tobacco Hornworm Tomato Hornworm	0.015 – 0.03	2.9 – 5.8	44 – 22	1	
	Beet Armyworm* Cabbage Looper Colorado Potato Beetle Cutworms Flea Beetle Grasshoppers Potato Aphid Southern Armyworm Tomato Fruitworm Tomato Pinworm Western Yellow-Striped Armyworm Whitefly	0.03 – 0.05	5.8 – 9.6	22 – 13		
	Vegetable Leafminer**	0.05	9.6	13		
	* Aids in control. ** Adjournal™ is not recommended for use on the Vegetable Leafminer in Florida. Do not apply more than 0.5 lbs. a.i. per acre per season.					
Turnips (TX only)	Armyworm Flea beetle Imported Cabbageworm	0.03 – 0.05	5.8 – 9.6	22 – 13	7	
	Do not apply more than 0.4 lbs a.i. per acre per season.					

**SPECIALTY USES**

Crop	Insect	Application Rate		Low Volume Sprays lb ai/acre
		High Volume Sprays		
		lb ai/100 gal	fl oz/100 gal	
Christmas tree plantings, Conifer plantations, Conifer seed orchards, Forest tree nurseries	Balsam Twig Aphid Balsam Woolly Adelgid Cranberry Girdler (adult control) European Pine Sawfly Nantucket Pine Tip Moth and other Pine Tip Moths (except CA) Pine Chafer Pine Conelet Bug Pine Needle Midge Pineleaf Chermid Red Pine Sawfly Redheaded Pine Sawfly Spittlebugs Spruce budworm	0.03 – 0.05	5.8 – 9.6	0.03 – 0.05
	Spray in sufficient gallonage to obtain good coverage of entire tree.			
	Northern Pine Weevil Pales Weevil	0.03 – 0.05	5.8 – 9.6	
	Dilute formulation in kerosene (rather than in water), and spray tree stumps to control developing larvae. Use sufficient spray volume to cover stump thoroughly.			
		Application Rate		
	Coneworm Seed Chalcid Seedbug	9.6 fl oz/100 gals of water for high volume sprayers. 52 fl oz/100 gals of water for low volume sprayers. 0.19 lb. ai/acre application in not less than 10 gals of water for aerial applications.		
Apply first application within 1 week of female flower closure or peak pollen flight for Webbing Coneworm control. For other Coneworms and Seedbugs, apply first application within 30 days following female flower closure. Repeat application at intervals of 4 weeks but do not apply more than 1.6 lbs. ai per acre per year. For Seed Chalcid control, apply when all cones are pendant, and repeat at 1 – 2 week intervals for 2 or more sprays. Apply approximately 5 – 10 gals of the 9.6 fl oz/100 gal dilution per tree with high volume sprayers. With low volume sprayers apply 100 gals of the 52 fl oz/100 gal dilution per acre. Do not graze or harvest cover crop. Refer to Spray Recommendations and Precautions when applying to areas adjacent to water.				

Crop	Insect	Application Rate		Acres treated per gal of Adjournal™
		lb ai/acre	fl oz/acre	
NON-CROPLAND (excluding public land such as forests, parks, or recreational)	Grasshoppers Saltmarsh Caterpillar	0.015 – 0.03	2.9 – 5.8	44 – 22
	Army Cutworms Armyworms Chinch Bugs	0.03 – 0.05	5.8 – 9.6	22 – 13
Spray non-cropland adjacent to tilled areas to control migrating insects (Grasshoppers, Armyworms) which are a threat to crops. Do not apply more than 0.5 lbs. active ingredient per acre per year. Do not feed treated crop to livestock. Refer to Spray Recommendations and Precautions when applying to areas adjacent to water.				

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store in a secure, dry and temperate area. Store in original container. Keep container closed when not in use. Do not store near food or feed. Do not use or store around the home. Avoid contact with water. In case of spill or leak, soak up with sand, earth or synthetic absorbent (do not use alkaline absorbents) and dispose of wastes in compliance with local, State and Federal regulations.

**PRODUCT DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent), then offer for recycling or reconditioning if container reuse is permitted. If container reuse is prohibited, puncture metal containers and dispose of in a sanitary landfill, or by other approved State and local procedures. Dispose of plastic containers in a sanitary landfill, or by incineration if allowed by State and local authorities. If burned, stay out of smoke.

In case of spill, avoid contact, isolate area, and keep out animals and unprotected persons. Confine spills. Call INFOTRAC: 1-800-535-5053.

**WARRANTY STATEMENT**

MAKHTESHIM AGAN OF NORTH AMERICA, INC. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of MAKHTESHIM AGAN OF NORTH AMERICA, INC. To the extent allowed by law, MAKHTESHIM AGAN OF NORTH AMERICA, INC. shall not be liable for consequential, special, or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. In addition to the foregoing, no purchaser of this product (other than an end user) shall be entitled to any reimbursement for any loss suffered as a result of any suspension or cancellation of the registration for this product by the U.S. Environmental Protection Agency. Except as expressly provided herein, MAKHTESHIM AGAN OF NORTH AMERICA, INC. makes no warranties, guarantees, or representations of any kind, either expressed or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. The exclusive remedy of any buyer or user of this product for any and all losses, injuries, or damages resulting from or in any way arising from the use, handling, or application of this product, whether in contract, warranty, tort, negligence, strict liability, or otherwise, shall be damages not exceeding the purchase price paid for this product or, at MAKHTESHIM AGAN OF NORTH AMERICA INC.'s election, the replacement of this product.

Adjournal™ is a trademark of Makhteshim Agan of North America, Inc.

DuPont™, "ASANA", "LANNATE" and "Super Tin" are trademarks or registered trademarks of E.I. duPont de Nemours and Company.

"Ambush", "Karate" are registered trademarks of Syngenta Crop Protection, Inc.

"Ammo", "Capture", "Fury", "Mustang", "Pounce" are registered trademarks of FMC Corp.

"Baythroid" is a registered trademark of Bayer CropScience.

"Dantol" is a registered trademark of Valent USA Corp.

"Decis", "Scout X-tra" are registered trademarks of Bayer CropScience SA.

(EPA app 01-05-05)(notif 07-17-06)