VIPER RTU

COUNTRY FRESH FRAGRANCE READY – TO – USE

For Domestic Indoor and Outdoor Use Only

Provides 60-Day (two months) Control of Fleas Provides 60-Day (two months) Control of Stored Product Pests

CONTROLS Fleas and Ticks on dogs, Carpet Beetles, Bed Bugs, Clothes Moth, Lice and Louse Eggs, Stable Flies, Horse Flies, Deer Flies, Aphids, Whiteflies, Cockroaches, Spiders, Red Mites and many more listed.

Read and use according to label directions. Store and transport in upright position.

EPA Reg. No. 4-393-53883

EPA Est. No. 53883-TX-002

ACTIVE INGREDIENT:	
ACTIVE INGREDIENT.	

Permethrin*	0.25%
INERT INGREDIENTS	
	$1\overline{00.00\%}$

*(3-phenoxyphenyl)methyl(+/-)cis/trans 3-(2,2 dichloroethenyl) 2,2-Dimethylcyclopropanecar-boxylate. Cis/trans ratio: min. 35% (+/-) cis and max. 65% (+/-) trans.

KEEP OUT OF REACH OF CHILDREN CAUTION

See inside booklet for additional Precautionary Statements

Net Contents: 1 Gal. (3.785 liters)

Manufactured For Control Solutions, Inc. 5903 Genoa-Red Bluff Pasadena, TX 77507-1041

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

Affix trigger, pump or other type sprayer to container for application. Do not transfer to another container.

SHAKE WELL BEFORE USING

INDOOR APPLICATION

For spot application only. Do not use as a space spray indoors.

SURFACE SPRAYING: FOR 60-DAY (TWO MONTHS) CONTROL OF: COCKROACHES, WATER-BUGS, PALMETTO BUGS, SCORPIONS, MILLIPEDES, CENTIPEDES, SOWBUGS, PILLBUGS, ANTS, SILVERFISH, FIREBRATS, SPIDERS, CRICKETS, CLOVER MITES, CHEESE MITES, GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, RUST RED FLOUR BEETLES, DRUGSTORE BEETLES, MEAL WORMS, GRAIN MITES, AND CADELLES: Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For *Silverfish*, spray bookcases. For *Ants*, spray trails, nests and points of entry.

TO CONTROL CARPET BEETLES: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

TO CONTROL BEDBUGS: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

TO CONTROL FLEAS (ADULTS AND LARVAE) AND TICKS (ADULTS AND LARVAE):

Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh beddings should be placed in animal quarters following treatment. Repeat treatment as needed. Allow to dry thoroughly before allowing pets to contact treated areas.

TO CONTROL MOTHS: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chest, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

TO KILL LICE AND LOUSE EGGS: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated particles to dry thoroughly before use.

ON LIVESTOCK: To protect cattle (beef and dairy), goats, sheep, hogs and horses from *Horn Flies*, *Face Flies*, *House Flies*, *Mosquitoes* and *Gnats*. Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

TO CONTROL STABLE FLIES, HORSE FLIES AND DEER FLIES: Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

TO CONTROL BLOOD-SUCKING LICE: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

TO CONTROL BEDBUGS AND MITES IN POULTRY HOUSES: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

ON DOGS

TO CONTROL FLEAS, TICKS, AND LICE: Start spraying at the tail, moving the dispenser rapidly and making sure that the animals entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly, bud do not saturate animal. Do not spray into eyes or face. Avoid contact with genitalia. Repeat as necessary.

NOTE: Not recommended for use on puppies less than four weeks old. DO NOT USE ON CATS OR KITTENS

OUTDOOR APPLICATION

To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6-10 feet wide and adjacent to activity areas, especially where dense vegetations occurs.

FOR CONTROL OF GARDEN INSECTS: For use on roses, dahlias, asters and other ornamentals to kill *Japanese Beetles* hit by spray as well as *Whiteflies, Aphids, Exposed Thrips, Spiders, Red Mites* and *Leafminers.* May be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, eunonymus, fuschia, geraniums, crassula, grapevines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed again Whiteflies, Aphids, Spider and Red Mites, Exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Armyworms.

When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals CAUTION

Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before surface applications. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwaters.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. THE MANUFACTURER MAKES NO WARRANTIES, EXPRESSED OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USE, WHICH EXTEND BEYOND THE DESCRIPTION ON THE LABEL. Read all directions carefully.